



# joint injection patient satisfaction survey: 2014


**Q1) I was examined by the Doctor Who did the injection before it was carried out**




**Q2) Other treatment options other than injection were explained to me.**




**Q3) I was told about possible side-effects or things that could go wrong with the injection.**



**Q4) I was satisfied that the doctor gave me all the information I needed before proceeding with the injection.**




**Q5) The doctor put me at ease doing the injection.**




**Q6) I was allowed to sit or lie in a comfortable position doing the injection.**







**Q7) The injection was not as sore as I thought it would be.**




**Q8) I was given clear instructions about resting the joint /muscle/tendon that had been injected.**




**Q09) I was given clear instructions about when to resume activity after the injection.**



**Q10) I was told what I could take if I had been at that injection.**



**Q11) I was clear about where I could get help if I had any problems after the injection.**



**Q12) I was satisfied with the arrangements that were made for me to be reviewed after the injection.**




Neutral	1	10%
Disagree	1	10%
Strongly disagree	0	0%
Not answered	0	0%
	<hr/>	
	10	




**Q13) After the injection I had increased pain.**

Not at all	5	50%
For 1 to 2 days	2	20%
For 3 to 5 days	1	10%
For 6 to 7 days	0	0%
More then one week	2	20%
Not answered	0	0%
	<hr/>	
	10	




**Q14) The injection helped relieve pain.**

Not at all	2	20%
For less than a week	0	0%
For between one week and one month	6	60%
For between one month and three months	1	10%
For between three months and six months	1	10%
For more than six months	0	0%
Not answered	0	0%
	<hr/>	
	10	




**Q15) The injection helped relieve stiffness**

Not at all	2	20%
For less than a week	2	20%
For between one week and one month	4	40%
For between one month and three months	1	10%
For between three months and six months	1	10%
For more than six months	0	0%
Not answered	0	0%
	<hr/>	
	10	



**Q16) The injection helped me to get back to "normal" activity.**

Not at all	2	20%
For less than a week	0	0%
For between one week and one month	6	60%
For between one month and three months	1	10%
For between three months and six months	1	10%
For more than six months	0	0%
Not answered	0	0%
	<hr/>	
	10	




**Q17) if the injection helped, would you say your problem was**


Completely cured	0	0%
A lot better	6	60%
A bit better	0	0%
Only a slight improvement	4	40%
Not answered	0	0%
	<hr/>	
	10	




**Q18) Overall I was satisfied with the outcome of the injection.**



**Q19) I would have the same thing done again if the problem recurred.**



Survey sent out	30
Survey returned	10



Comments received:

- 1) I feel better
- 2) I had no complications.
- 3) Very grateful for the hand injections, but the injection clinics can be a little longer to sort out. This can make my hands sore as I have injections into each hand at the best every three months, but understandably the clinics may not run that frequent.
- 4) My knees have been painful for years. I normally descend the stairs right foot first. After the injection I was feeling comfortable enough to walk down the steps normally. However, after two weeks the pain was severe causing my stride change.